

PLANNING YOUR REVUFORJ ROUTINE

A medication tracker for patients and caregivers

Taking Revuforj Tablets: Information for you or your loved one

It's important to take Revuforj exactly as your healthcare provider tells you to. Do not change your dose or stop taking Revuforj unless your healthcare provider tells you to.

Your healthcare provider is the best source of information when it comes to your health.



Bottles and tablets shown are not actual size.

What is Revuforj?

Revuforj® (revumenib) is a prescription medicine used to treat adults and children 1 year and older with acute leukemia with a lysine methyltransferase 2A gene (*KMT2A*) translocation whose disease has come back or has not improved after previous treatment(s).

Your healthcare provider will perform a test to make sure that Revuforj is right for you. It is not known if Revuforj is safe and effective in children less than 1 year of age.

Important Safety Information

What is the most important information I should know about Revuforj?

Revuforj may cause serious side effects, including: Differentiation syndrome. Differentiation syndrome is a serious but common condition that affects your blood cells, which may be life-threatening or lead to death if not treated. Differentiation syndrome has happened as early as 3 days and up to 41 days after starting Revuforj. Tell any healthcare provider caring for you that you are taking a medicine that can cause differentiation syndrome. Call your healthcare provider or go to the nearest hospital emergency room right away if you develop any of the following symptoms of differentiation syndrome while taking Revuforj:

- fever
- cough
- shortness of breath
- severe headache
- confusion
- dizziness or lightheadedness
- fast weight gain
- swelling of arms, legs, neck, groin, or underarm area
- decreased urination

If you develop any of these symptoms of differentiation syndrome, your healthcare provider may start you on a medicine given through a vein (intravenous) called corticosteroids and may monitor you in the hospital.

Please see additional Important Safety Information on page 4 and Full Prescribing Information, including **BOXED WARNING**, and Medication Guide and Instructions for Use.

 **Revuforj**[®]
(revumenib) tablets
25 mg • 110 mg • 160 mg

HOW TO TAKE REVUFORJ TABLETS

Revuforj tablets come in different strengths. Each strength is a different color:



Tablets shown are not actual size.

Your healthcare provider may prescribe more than 1 strength of Revuforj tablets for you, so it is important that you understand how to take your medicine the right way. Be sure that you understand exactly how many tablets you need to take, and what strengths to take.

All Revuforj doses should be taken:



2 times a day by mouth

Take at about the same time each day (about 12 hours apart)



On an empty stomach

In the clinical trial, this was defined as at least 2 hours after a meal and 1 hour before the next meal

OR



With a low-fat meal

Low-fat meals should be about 400 calories and contain 25% or less fat

- Swallow Revuforj tablets whole with a cup of water
- **Do not** cut or chew tablets
- If you are unable to swallow tablets, Revuforj can be crushed and dispersed in water as directed in the **Instructions for Use**.
 - See the **Instructions for Use** for detailed information on how to prepare and give Revuforj



If you miss a dose of Revuforj or did not take it at the usual time, take your dose as soon as possible and at least 12 hours before your next dose. **Do not** take 2 doses within 12 hours. Return to your normal scheduled dose the following day.

- Store Revuforj at room temperature between 68°F to 77°F (20°C to 25°C)
- Keep the tablets in the bottle that it comes in until you are ready to take it
- The Revuforj bottle has a drying agent (desiccant) and child-resistant closure

Keep Revuforj and all medicines out of reach of children

Please see Important Safety Information on pages 1 and 4 and **Full Prescribing Information**, including **BOXED WARNING**, and **Medication Guide and Instructions for Use**.

GETTING INTO A REVUFORJ ROUTINE




































Planning a routine right from the start can be a helpful way to make sure Revuforj is taken as directed. Here are some tips that may help you get started:

- Keep Revuforj somewhere you access every day, but safely away from children
- Set an alarm on your phone or leave a reminder note in plain sight
- Plan to take Revuforj at the same time each day. For example, 7 AM and 7 PM

Use the example calendar below to help track your Revuforj dosing schedule

Fill in the **month** and your **prescribed dose** of the tablets. Use a current calendar to fill in the dates for each day. **Check off** or **cross out** each morning and evening dose right after you take it.

MONTH: _____ YOUR DOSE: _____ mg

SUN	M	T	W	TH	F	SAT
						
						
						
						
						

All doses are taken 2 times a day on an empty stomach or with a low-fat meal.
Always check with your doctor for your exact dose and schedule.

See the last page to print more calendars to continue to track your treatment.

 **Revuforj**[®]
(revumenib) tablets
25 mg • 110 mg • 160 mg

STAY TRACK

For more tips on how to stay organized and plan a routine, visit [Revuforj.com](https://www.Revuforj.com)

Important Safety Information (cont'd)

Changes in electrical activity of your heart called QT prolongation. QT prolongation is a serious but common side effect that can cause irregular heartbeats that can be life-threatening or lead to death. Your healthcare provider will check the electrical activity of your heart with a test called an electrocardiogram (ECG) and will also do blood tests to check your potassium and magnesium levels before and during treatment with Revuforj. Tell your healthcare provider right away if you feel faint, lightheaded, dizzy, or if you feel your heart beating irregularly or fast during treatment with Revuforj.

Before taking Revuforj, tell your healthcare provider about all your medical conditions, including if you:

- have any heart problems, including a condition called long QT syndrome
- have been told you have low blood levels of potassium or magnesium
- are pregnant or plan to become pregnant. Revuforj can harm your unborn baby
 - Your healthcare provider will perform a pregnancy test within 7 days before you start treatment with Revuforj. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with Revuforj
 - **Females** who are able to become pregnant should use effective birth control (contraception) during treatment with Revuforj and for 4 months after the last dose of Revuforj
 - **Males** who have female partners who are able to become pregnant should use effective birth control during treatment with Revuforj and for 4 months after the last dose of Revuforj
 - Talk to your healthcare provider about birth control methods you can use during this time
 - Revuforj may cause fertility problems in females and males. Talk to your healthcare provider if this is a concern for you
- are breastfeeding or plan to breastfeed. It is not known if Revuforj passes into your breast milk. Do not breastfeed during your treatment with Revuforj or for 1 week after your last dose of Revuforj

Tell your healthcare provider about all the medications you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Revuforj and other medicines may affect each other causing side effects.

The most common side effects of Revuforj include:

- bleeding (hemorrhage)
- low white blood cell counts with fever
- swelling in the arms and legs
- nausea and vomiting
- decreased appetite
- muscle pain
- diarrhea
- constipation
- infections, including bacterial and viral
- changes in liver function tests
- tiredness

Your healthcare provider will do blood tests and ECGs before you start and during treatment with Revuforj. Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with Revuforj if you develop certain side effects. These are not all the possible side effects of Revuforj. Call your healthcare provider for medical advice about side effects.

You are encouraged to report side effects of prescription drugs to FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.




































Please see [Full Prescribing Information](#), including **BOXED WARNING, and [Medication Guide and Instructions for Use](#).**

STAY TRACK

Use the calendar below to help track your dosing schedule

Fill in the **month** and your **prescribed dose** of tablets. Use a current calendar to fill in the dates for each day. **Check off** or **cross out** each morning dose and evening dose right after you take it.

MONTH: _____ YOUR DOSE: _____ mg

SUN	M	T	W	TH	F	SAT
						
						
						
						
						

All doses are taken 2 times a day on an empty stomach or with a low-fat meal.
Always check with your doctor for your exact dose and schedule.